

# Cosmic Censorship

## in a Gravitational Collapse in Quantum Einstein Gravity

Benjamin Koch

in colaboraion with  
Alfio Bonanno and Alessia Platania  
bkoch@fis.puc.cl

<sup>a</sup> PUC, Chile

Leiden  
February 2017




- Motivation
- Black holes in Asymptotic Safety
- Cosmic Censorship in Asymptotic Safety
- Conclusion

[\*]Based on: A. Bonanno, B.K., A. Platania, e-Print: arXiv:1610.05299



# Motivation

FRGE solutions:



[\*]M. Reuter, F. Saueressig, Phys.Rev. D65 (2002) 065016; probably also all people present...


- What does this mean for physical systems?

e.g. BLACK HOLES



# Motivation

FRGE solutions:



[\*]M. Reuter, F. Saueressig, Phys.Rev. D65 (2002) 065016; probably also all people present...

- What does this mean for physical systems?


e.g. **BLACK HOLES**



# Black Holes



# Black Holes




Black Holes



# Black Holes



Singularitiy




Black Holes




# Black Holes



Singularity



Asymptotic Safety?



Black Holes

Black holes in Asymptotic Safety







# Black Holes



Singularity



Asymptotic Safety?



Black Holes

Black holes in Asymptotic Safety



Black holes in Asymptotic Safety:  
Two approaches borrowed from QFT

- Improving solutions (Uehling potential textbook QED)
- Improving action and eom (gap equations in QFT)



# Black Holes

## Improving solutions:

Classical eom's

$$G_{\mu\nu} + g_{\mu\nu}\Lambda = 8\pi G_k T_{\mu\nu} \quad (1)$$

Classical solution for  $ds^2 = f(r)dt^2 + f^{-1}dr^2 + d\Omega$  (with  $\Lambda_k \approx 0$ )

$$f(r) = 1 - \frac{2G_k M}{r} \quad (2)$$

Quantum improvement  $G_k$  with  $k \neq \text{cte.}$

$$k = k(r) = \frac{\xi}{d(r)} \quad (3)$$


where  $d(r)$  physical cut-off like proper distance \*

[\*] A. Bonanno, M. Reuter, Phys. Rev. D62, 043008 (2000)



# Black Holes

## Improving solutions:\*



- No Singularity
- Stable remnant
- Similar for different scale setting, extra dimensions, charge, or angular momentum **but**

[\*] A. Bonanno, M. Reuter Phys.Rev. D62 (2000) 043008; figure from B.K., F. Saueressig, Int.J.Mod.Phys. A29 (2014) no.8, 1430011



# Black Holes

Improving solutions:\*


but if one considers

$$\Lambda_k|_{UV} = \lim_{k \rightarrow \infty} k^2 \lambda^* \quad (4)$$

$\Rightarrow$  the neglected term  $\sim \Lambda_k$  in lapse function

$$f(r) = 1 - \frac{2G_k M}{r} + r^2 \Lambda_k \quad (5)$$

can become **divergent** for  $r \rightarrow 0_*$



[\*] B.K., F. Saueressig, Class.Quant.Grav. 31 (2014) 015006



# Black Holes

## Gap equations:\*

Effective Einstein-Hilbert action

$$\Gamma_k[g_{\mu\nu}] = \int_M d^4x \sqrt{-g} \left( \frac{R - 2\Lambda_k}{16\pi G_k} \right) , \quad (6)$$

eom  $\delta g_{\mu\nu}$ :

$$G_{\mu\nu} = -g_{\mu\nu}\Lambda_k - \Delta t_{\mu\nu} + 8\pi G_k T_{\mu\nu} , \quad (7)$$

with

$$\Delta t_{\mu\nu} = G_k (g_{\mu\nu} \square - \nabla_\mu \nabla_\nu) \frac{1}{G_k} . \quad (8)$$

scale setting  $\delta k$ :\*

$$\left[ R \partial_k \left( \frac{1}{G_k} \right) - 2 \partial_k \left( \frac{\Lambda_k}{G_k} \right) \right] = 0$$

[\*] B.K., P. Riaseco, C. Contreras Phys.Rev. D91 (2015) no.2, 025009



# Black Holes

## Gap equations:

Complicated equations  $\Rightarrow$  no analytic BH solution

Trick: Impose Null Energy Condition

$$\nabla_{\mu} \Delta t^{\mu\nu} = 0 \quad (10)$$

Trick implies Schwarzschild ansatz  $g_{00} = 1/g_{11} = f(r)$

$\Rightarrow$  generalized de Sitter solution, also Reissner Nordstrom, and BTZ:[\*]

$$G(r) = \frac{G_0}{\epsilon r + 1} \quad (11)$$

$$f(r) = 1 + 3G_0 M_0 \epsilon - \frac{2G_0 M_0}{r} - (1 + 6\epsilon G_0 M_0) \epsilon r - \frac{\Lambda_0 r^2}{3} + r^2 \epsilon^2 (6\epsilon G_0 M_0 + 1) \ln \left( \frac{c_4(\epsilon r + 1)}{r} \right) \quad (12)$$

...

Constants of integration:  $G_0$ ,  $M_0$ ,  $\Lambda_0$ ,  $\epsilon$ ,  $c_4$


[\*] B.K., P. Rioseco, Class.Quant.Grav. 33 (2016) 035002,

B.K. I. Reyes, A. Rincon, Class.Quant.Grav. 33 (2016) no.22, 225010.



# Black Holes

## Gap equations:



- Has singularity ...





Fair to say:

Question of singularity is still open!

What is the problem with such singularities?



Fair to say:

Question of singularity is still open!

What is the problem with such singularities?




# Black Hole Formation



# Black Hole Formation

Remember classical BH



Black Holes




# Black Hole Formation

Remember classical BH



Singularitiy



Black Holes



# Black Hole Formation

Remember classical BH



Singularities



Censorship hypothesis



Black Holes

**dressed** singularity might not be the problem

⇒

study **naked** singularities (e.g. BH formation)



# Black Hole Formation

Remember classical BH



Singularities



Censorship hypothesis




Black Holes

**dressed** singularity might not be the problem  
⇒  
study **naked** singularities (e.g. BH formation)



# Black Hole Formation

Classical Kuroda-Papapetrou model



Black Hole formation





# Black Hole Formation

Classical Kuroda-Papapetrou model



Singularity



Black Hole formation



# Black Hole Formation

Classical Kuroda-Papapetrou model



Singularity



Censorship hypothesis  
comes "late"



Black Hole formation




# Black Hole Formation

Classical Kuroda-Papapetrou model



Singularity



Censorship “late”, AS  
can help?



Black Hole formation



# Black Hole Formation

## Classical Kuroda-Papapetrou model

Classical Vaidya metric

$$ds^2 = -f(r, v) \cdot dv^2 + 2dvdr + r^2 d\Omega^2 \quad (13)$$

with advanced ingoing null coordinate  $v$ .

Null geodesics:

$$\frac{dr}{dv} = \frac{1}{2} \left( 1 - \frac{2G_0 m(v)}{r} \right). \quad (14)$$

$$f(r, v) = 1 - \frac{2G_0 m(v)}{r} \quad (15)$$

Mass inflow modeled by:

$$m(v) = \begin{cases} 0 & v < 0 \\ \lambda v & 0 \leq v < \bar{v} \\ \bar{m} & v \geq \bar{v} \end{cases}$$



# Black Hole Formation

## Classical Kuroda-Papapetrou model

Horizons:

- High mass inflow

$$\lambda > \lambda_c = \frac{1}{16G_0} \quad (17)$$

⇒ Singularity at  $r = 0$  **always** covered by an horizon

- Low mass inflow

$$\lambda < \lambda_c = \frac{1}{16G_0} \quad (18)$$

⇒ Singularity at  $r = 0$  can be **naked**


Can be seen in phase diagram:



# Black Hole Formation

Classical Kuroda-Papapetrou model


Phase diagram:



Singularity covered

blue apparent horizon, purple event horizon

Singularity naked



# Black Hole Formation

AS improved Kuroda-Papapetrou model

Improved Vaidya metric

$$ds^2 = -f_k(r, v) \cdot dv^2 + 2dvdr + r^2 d\Omega^2 \quad (19)$$

with

$$f_k(r, v) = 1 - \frac{2G_k m(v)}{r} \quad (20)$$

Identify IR cut-off scale with scale imposed by infalling radiation

$$k \sim T \sim \rho^{1/4} \quad (21)$$

$\xi$  : proportionality constant,  $\rho$  given from classical field equations ( $G_v, v$ )

$$\frac{\dot{m}(v)}{4\pi r^2} = \rho(v, r). \quad (22)$$

Thus,

$$f_k(r, v) = 1 - \frac{2\lambda G_0 v}{r + \alpha\sqrt{\lambda}}, \quad \text{with } \alpha = \frac{\xi^2 G_0}{\sqrt{4\pi g_*}}$$



# Black Hole Formation

## AS improved Kuroda-Papapetrou model

Note:[\*]

- Improved lapse function  $f_k(r, v)$  is well defined in the limit  $r \rightarrow 0$

$$\lim_{r \rightarrow 0} f_k(r, v) = 1 - \frac{\sqrt{16\pi\lambda}}{\omega \xi^2} v \quad (24)$$

- However **singular** curvatures in  $r \rightarrow 0$  e.g.

$$R = -\frac{G_0\sqrt{\lambda}v}{\alpha r^2} + O(1/r^2), \quad K = \frac{16G_0\sqrt{\lambda}v}{\alpha^2 r^4} + O(1/r^3).$$

- One might invent cut-off identification **without** singularity, but don't want to do reverse engineering
- Like in all improving solutions schemes (24) does not solve eoms

[\*] B. Bonanno, B.K., A. Platania, arXiv:1610.05299.





# Black Hole Formation

AS improved Kuroda-Papapetrou model

From (24) apparent horizon shifted by the constant  $\alpha \sqrt{\lambda}$

$$r_{\text{AH}}(v) = 2 m(v) G_0 - \alpha \sqrt{\lambda} = 2 m(v) G_0 - \frac{G_0 \xi^2}{g_*} \sqrt{\frac{\lambda}{4\pi}}, \quad (25)$$

from  $r_{\text{AH}} \geq 0$  and matching to improved Schwarzschild  $\rightarrow$  minimum “time”  $\bar{v}$  of irradiation, necessary to actually form a black hole

$$r_S = 2 \lambda \bar{v} G_0 - \alpha \sqrt{\lambda} \geq 0 \quad \Rightarrow \quad \bar{v} \geq v_{\text{min}}(\lambda) \equiv \frac{\xi^2}{2 g_*} \sqrt{\frac{1}{4\pi\lambda}}. \quad (26)$$



# Black Hole Formation

AS improved Kuroda-Papapetrou model

Null geodesics from

$$\dot{r}(v) = \frac{1}{2} \left( 1 - \frac{2\lambda v G_0}{r(v) + \alpha\sqrt{\lambda}} \right), \quad (27)$$

Integrating (e.g. for  $\lambda \leq \frac{1}{16G_0}$ ) gives implicit equation

$$\frac{|r(v) + \alpha\sqrt{\lambda} - \mu_- v|^{\mu_-}}{|r(v) + \alpha\sqrt{\lambda} - \mu_+ v|^{\mu_+}} = \tilde{C} \quad (28)$$

with two linear solutions


$$r_{\pm}(v) = -\alpha\sqrt{\lambda} + \mu_{\pm} v,$$



# Black Hole Formation


AS improved Kuroda-Papapetrou model

Phase diagram:



blue apparent horizon, purple event horizon


Singularity **always naked** but how bad is it?




# Black Hole Formation

AS improved Kuroda-Papapetrou model

Phase diagram:



blue apparent horizon, purple event horizon  
Singularity **always naked** but how bad is it?



# Black Hole Formation

## AS improved Kuroda-Papapetrou model

Nature of the singularity (how bad is it?)

Study geodesics as dynamical system<sup>[\*]</sup>

$$\begin{cases} \frac{dv(t)}{dt} = N(r, v) \\ \frac{dr(t)}{dt} = D(r, v) \end{cases}, \quad (30)$$

where  $t$  is a parameter and the functions  $N(r, v)$  and  $D(r, v)$  are defined as

$$N(r, v) = 2r \quad D(r, v) = r - 2M(r, v). \quad (31)$$

Singularities are fixed points (e.g.  $r = 0$  and  $M(0, v) = 0$ )

Expand near the singularity

$$\begin{cases} \frac{dv(t)}{dt} = \dot{N}_{\text{FP}} (v - v_{\text{FP}}) + N'_{\text{FP}} (r - r_{\text{FP}}) \\ \frac{dr(t)}{dt} = \dot{D}_{\text{FP}} (v - v_{\text{FP}}) + D'_{\text{FP}} (r - r_{\text{FP}}) \end{cases}.$$

[\*] M. D. Mkenyeleze, R. Goswami, and S. D. Maharaj, Phys. Rev. D 90, 064034 (2014).



# Black Hole Formation

AS improved Kuroda-Papapetrou model

Nature of the singularity classified by eigenvalues of the stability matrix  $J$  of the system (32)

$$\chi_{\pm} = \frac{1}{2} \left( \text{Tr}J \pm \sqrt{(\text{Tr}J)^2 - 4 \det J} \right), \quad (33)$$

where

$$\text{Tr}J = \dot{N}_{\text{FP}} + D'_{\text{FP}} = 1 - 2 M'_{\text{FP}} \quad (34)$$

$$\det J = \dot{N}_{\text{FP}} D'_{\text{FP}} - \dot{D}_{\text{FP}} N'_{\text{FP}} = 4 \dot{M}_{\text{FP}}. \quad (35)$$



# Black Hole Formation

AS improved Kuroda-Papapetrou model

Strength of the singularity is

$$S = \frac{\dot{M}_{FP} X_{FP}^2}{2} = 0. \quad (36)$$

where  $X_{FP} \equiv \lim_{(r,v) \rightarrow FP} \frac{v(r)}{r}$ .

$\Rightarrow$  singularity is **integrable** "harmless".

Interesting:

- $S \rightarrow 0$  does not depend on cut-off identification as long as

$$\lim_{r \rightarrow 0} G_k(r) = \lim_{k \rightarrow \infty} G_k = 0$$



# Summary






# Summary

- Quantum gravity and Asymptotic Safety
- Black holes in AS: singularity unsure
- Naked singularities e.g. Kuroda-Papapetrou model
- AS improved Kuroda-Papapetrou model



# Summary

Take home messages:



- Important test QG candidate with problematic solutions of GR
- In different attempts, the singularity might go away or persist
- Even if naked singularities don't go away in AS, at least they become integrable



# Thank you

Thank you !

